Brain Research, 593 (1992) 347–349
© 1992 Elsevier Science Publishers B.V. All rights reserved 0006-8993/92/$05.00

Author Index

Abe, K., Ishiyama, J. and Saito, H.
Effects of epidermal growth factor and basic fibroblast growth factor on generation of long-term potentiation in the dentate gyrus of fimbria-fornix-lesioned rats (593) 335
Abe, K., see Chida, N. (593) 57
Akiyama, K., see Kohira, I. (593) 82
Altemus, M., Cizza, G. and Gold, P.W.
Chronic fluoxetine treatment reduces hypothalamic vasopressin secretion in vitro (593) 311
Aoki, C., see Farb, C. (593) 145

Balasubramaniam, A., see Chance, W.T. (593) 20
Barna, I. and Koenig, J.I.
Effects of mediobasal hypothalamic lesion on immunoreactive ACTH/β-endorphin levels in cerebrospinal fluid, in discrete brain regions, in plasma, and in pituitary of the rat (593) 69
Bates, C.M., see Haun, S.E. (593) 45
Bennett-Clarke, C., see Shen, E. (593) 136
Bernardi, G., see Mercuri, N.B. (593) 51
Besson, J.M., see Kayser, V. (593) 7
Beyer, C., Kolbinger, W., Froehlich, U., Pilgrim, C. and Reisert, I.
Sex differences of hypothalamic prolactin cells develop independently of the presence of sex steroids (593) 253
Bockaert, J., see Lafon-Cazal, M. (593) 63
Bougault, I., see Lafon-Cazal, M. (593) 63
Bush, L.G., see Midgley, L.P. (593) 89
Büsselberg, D., Michael, D., Evans, M.L., Carpenter, D.O. and Haas, H.L.
Zinc (Zn^{2+}) blocks voltage gated calcium channels in cultured rat dorsal root ganglion cells (593) 77

Calabresi, P., see Mercuri, N.B. (593) 51
Cammarata, S., see Tabaton, M. (593) 299
Caramia, M.D., see Rossini, P.M. (593) 14
Carpenter, D.O., see Büsselberg, D. (593) 77
Carter, R.E., see Meyerhoff, J.L. (593) 140
Cavazos, J.E., see Golarai, G. (593) 257
Chance, W.T., Balasubramaniam, A., Thomas, I. and Fischer, J.E.
Amylin increases transport of tyrosine and tryptophan into the brain (593) 20
Chen, J., see Meldrum, B.S. (593) 1
Chida, N., Saito, H. and Abe, K.
Spermine facilitates the generation of long-term potentiation of evoked potential in the dentate gyrus of anesthetized rats (593) 57
Cizza, G., see Altemus, M. (593) 311
Clemens, J.A., see Stephenson, D.T. (593) 128
Condés-Lara, M., see Hernández, J.R. (593) 239
Costentin, J., see Naudon, L. (593) 32
Côté, P.-Y. and Parent, A.
Calbindin and choline acetyltransferase are expressed by different neuronal populations in pedunculopontine nucleus but not in nucleus basalis in squirrel monkeys (593) 245
Coyle, J.T., see Meyerhoff, J.L. (593) 140

Daigen, A., see Kohira, I. (593) 82
D'Amelio, F., see Hyde, T.M. (593) 291
Daunton, N.G., see Hyde, T.M. (593) 291
Desiato, M.T., see Rossini, P.M. (593) 14
Doubleday, B. and Robinson, P.P.
The role of nerve growth factor in collateral reinnervation by cutaneous C-fibres in the rat (593) 179
Dreifuss, J.J., see Widmer, H. (593) 215
Dun, N.J., see Shen, E. (593) 136
Dun, S.L., see Shen, E. (593) 136
Dunwiddie, T.V., see Lupica, C.R. (593) 226

Evans, M.L., see Büsselberg, D. (593) 77

Farb, C., Aoki, C., Milner, T., Kaneko, T. and LeDoux, J.
Glutamate immunoreactive terminals in the lateral amygdaloid nucleus: a possible substrate for emotional memory (593) 145
Fercakova, A., Halat, G., Marsala, M., Lukacova, N. and Marsala, J.
Graded postischemic reoxygenation reduces lipid peroxidation and reperfusion injury in the rabbit spinal cord (593) 159
Fischer, J.E., see Chance, W.T. (593) 20
Foley, T.D. and Rhoads, D.E.
Effects of ethanol on Na^+-dependent amino acid uptake: dependence on rat age and Na^+,K^+-ATPase activity (593) 39
Froehlich, U., see Beyer, C. (593) 253

Gale, K., see Pasini, A. (593) 287
Gambetti, P., see Tabaton, M. (593) 299
Ge, N., see Masliah, E. (593) 323
Getchell, M.L., see Krishna, N.S.R. (593) 295
Getchell, T.V., see Krishna, N.S.R. (593) 295
Gibb, J.W., see Midgley, L.P. (593) 89
Golarai, G., Cavazos, J.E. and Sutula, T.P.
Activation of the dentate gyrus by pentylenetetrazol evoked seizures induces mossy fiber synaptic reorganization (593) 257
Gold, P.W., see Altemus, M. (593) 311
Good, P.F., Olanow, C.W. and Perl, D.P.
Neuromelanin-containing neurons of the substantia nigra accumulate iron and aluminum in Parkinson's disease: a LAMMA study (593) 343
Graham, S.H., see Meldrum, B.S. (593) 1
Guilbaud, G., see Kayser, V. (593) 7
Gwinn, R., see Meldrum, B.S. (593) 1

Haas, H.L., see Büsselberg, D. (593) 77
Halat, G., see Fercakova, A. (593) 159
Hammer Jr., R.R., see Hohmann, A.G. (593) 265
Hanai, K., see Tooyama, I. (593) 274
Hanna, R.B., see Marcus, D.S. (593) 185
Hanson, G.R., see Midgley, L.P. (593) 89
Haun, S.E., Murphy, E.J., Bates, C.M. and Horrocks, L.A.
Extracellular calcium is a mediator of astroglial injury during combined glucose-oxygen deprivation (593) 45
Hemstreet, M.K., see Hohmann, A.G. (593) 265
Hernández, J.R. and Condés-Lara, M.
Brain Na^+/K^+-ATPase regulation by serotonin and norepinephrine in normal and kindled rats (593) 239
Hohmann, A.G., Matsumoto, R.R., Hemstreet, M.K., Patrick, S.L., Margulies, J.E., Hammer Jr., R.R. and Walker, J.M.
Effects of 1,3-di-*o*-tolylguanidine (DTG), a sigma ligand, on local cerebral glucose utilization in rat brain (593) 265
Hong, J.S., see McMillian, M. (593) 307
Horrocks, L.A., see Haun, S.E. (593) 45
Hyde, T.M., Wu, L.-C., Krasnov, I.B., Sigworth, S.K., Daunton, N.G. and D'Amelio, F.
Quantitative autoradiographic analysis of muscarinic cholinergic and $GABA_A$ (benzodiazepine) receptors in the forebrain of rats flown on the Soviet Biosatellite COSMOS 2044 (593) 291

Ishikawa, E., see Ohgo, S. (593) 25
Ishiyama, J., see Abe, K. (593) 335
Isom, G.E., see Patel, M.N. (593) 114

Jensen, R.T., see Ladenheim, E.E. (593) 168

Kadota, K., see Meldrum, B.S. (593) 1
Kaneko, T., see Farb, C. (593) 145
Kawai, M., see Tabaton, M. (593) 299
Kayser, V., Guilbaud, G. and Besson, J.M.
Potent antinociceptive effects of clonidine systemically administered in an experimental model of clinical pain, the arthritic rat (593) 7
Kimura, H., see Tooyama, I. (593) 274
Kisara, K., see Sakurada, T. (593) 319
Koenig, J.I., see Barna, I. (593) 69
Kohira, I., Akiyama, K., Daigen, A. and Otsuki, S.
Enduring increase in membrane-associated protein kinase C activity in the hippocampal-kindled rat (593) 82
Kolbinger, W., see Beyer, C. (593) 253
Koob, G.F., see Weiss, F. (593) 314
Krasnov, I.B., see Hyde, T.M. (593) 291
Kriebel, M.E., see Marcus, D.S. (593) 185

Krishna, N.S.R., Getchell, T.V., Margolis, F.L. and Getchell, M.L.
Amphibian olfactory receptor neurons express olfactory marker protein (593) 295
Kuzmin, A., Patkina, N., Pchelintsev, M. and Zvartau, E.
Isradipine is able to separate morphine-induced analgesia and place conditioning (593) 221

Ladenheim, E.E., Jensen, R.T., Mantey, S.A. and Moran, T.H.
Distinct distributions of two bombesin receptor subtypes in the rat central nervous system (593) 168
Lafon-Cazal, M., Bougault, I., Steinberg, R., Pin, J.P. and Bockaert, J.
Measurement of γ-enolase release, a new method for selective quantification of neurotoxicity independently from glial lysis (593) 63
Lampis, G., see Olianas, M.C. (593) 304
Leach, M.J., see Meldrum, B.S. (593) 1
LeDoux, J., see Farb, C. (593) 145
Leroux-Nicollet, I., see Naudon, L. (593) 32
Levy, A.D., see Rittenhouse, P.A. (593) 105
Levy, W.B., see Zhang, D.X. (593) 124
Li, Q., see Rittenhouse, P.A. (593) 105
Lorang, M.T., see Weiss, F. (593) 314
Lukacova, N., see Fercakova, A. (593) 159
Lupica, C.R., Proctor, W.R. and Dunwiddie, T.V.
Dissociation of μ and δ opioid receptor-mediated reductions in evoked and spontaneous synaptic inhibition in the rat hippocampus in vitro (593) 226

Mallory, M., see Masliah, E. (593) 323
Mandybur, T., see Tabaton, M. (593) 299
Manome, Y., see Sakurada, T. (593) 319
Mantey, S.A., see Ladenheim, E.E. (593) 168
Marcus, D.S., Kriebel, M.E. and Hanna, R.B.
Effects of calcium on the dynamic process of transmitter release which generates either skew- or bell-MEPPS (593) 185
Margolis, F.L., see Krishna, N.S.R. (593) 295
Margulies, J.E., see Hohmann, A.G. (593) 265
Markou, A., see Weiss, F. (593) 314
Marsala, J., see Fercakova, A. (593) 159
Marsala, M., see Fercakova, A. (593) 159
Masliah, E., Mallory, M., Ge, N. and Saitoh, T.
Amyloid precursor protein is localized in growing neurites of neonatal rat brain (593) 323
Matsukura, S., see Ohgo, S. (593) 25
Matsumoto, R.R., see Hohmann, A.G. (593) 265
McGeer, E.G., see Tooyama, I. (593) 274
McGeer, P.L., see Shigematsu, K. (593) 117
McGeer, P.L., see Tooyama, I. (593) 274
McMillian, M., Hong, J.S. and Pennypacker, K.R.
Preferential activation of [^{3}H]phorbol-12,13-dibutyrate binding by AMPA (α-amino-3-hydroxy-5-methyl-4-isoxazole-propionic acid) in neonatal striatal cell cultures (593) 307
Meldrum, B.S., Swan, J.H., Leach, M.J., Millan, M.H., Gwinn, R., Kadota, K., Graham, S.H., Chen, J. and Simon, R.P.
Reduction of glutamate release and protection against ischemic brain damage by BW 1003C87 (593) 1
Mercuri, N.B., Stratta, F., Calabresi, P. and Bernardi, G.
A voltage-clamp analysis of NMDA-induced responses on dopaminergic neurons of the rat substantia nigra zona compacta and ventral tegmental area (593) 51
Meyerhoff, J.L., Carter, R.E., Yourick, D.L., Stauch Slusher, B. and Coyle, J.T.
Genetically epilepsy-prone rats have increased brain regional activity of an enzyme which liberates glutamate from *N*-acetyl-aspartyl-glutamate (593) 140
Michael, D., see Büsselberg, D. (593) 77
Midgley, L.P., Bush, L.G., Gibb, J.W. and Hanson, G.R.
Characterization of phencyclidine-induced effects on neuropeptide Y systems in the rat caudate-putamen (593) 89
Millan, M.H., see Meldrum, B.S. (593) 1
Milner, T., see Farb, C. (593) 145
Moran, T.H., see Ladenheim, E.E. (593) 168
Mori, N. and Ohta, S.-i.
Comparison of anticonvulsant effects of valproic acid entrapped in positively and negatively charged liposomes in amygdaloid-kindled rats (593) 329
Murphy, E.J., see Haun, S.E. (593) 45

Nakatsuru, K., see Ohgo, S. (593) 25
Naudon, L., Leroux-Nicollet, I. and Costentin, J.
Consequences of an intrastriatal injection of kainic acid on the dopaminergic neuronal and vesicular uptake systems (593) 32

Ohba, M., see Sakurada, T. (593) 319
Ohgo, S., Nakatsuru, K., Ishikawa, E. and Matsukura, S.
Stimulation of cholecystokinin (CCK) release from superfused rat hypothalamo–neurohypophyseal complexes by interleukin-1 (IL-1) (593) 25
Ohta, S.-i., see Mori, N. (593) 329
Olanow, C.W., see Good, P.F. (593) 343
Olianas, M.C., Lampis, G. and Onali, P.
Human Y-79 retinoblastoma cells express functional corticotropin-releasing hormone receptors (593) 304
Onali, P., see Olianas, M.C. (593) 304
Otsuki, S., see Kohira, I. (593) 82

Parent, A., see Côté, P.-Y. (593) 245
Pasini, A., Tortorella, A. and Gale, K.
Anticonvulsant effect of intranigral fluoxetine (593) 287
Patel, M.N., Yim, G.K.W. and Isom, G.E.
Potentiation of cyanide neurotoxicity by blockade of ATP-sensitive potassium channels (593) 114
Patkina, N., see Kuzmin, A. (593) 221
Patrick, S.L., see Hohmann, A.G. (593) 265
Pchelintsev, M., see Kuzmin, A. (593) 221
Pennypacker, K.R., see McMillian, M. (593) 307
Perl, D.P., see Good, P.F. (593) 343
Perry, G., see Tabaton, M. (593) 299
Peterson, M.R., see Robertson, H.A. (593) 332
Pilgrim, C., see Beyer, C. (593) 253
Pin, J.P., see Lafon-Cazal, M. (593) 63
Proctor, W.R., see Lupica, C.R. (593) 226

Raggenbass, M., see Widmer, H. (593) 215
Rash, K., see Stephenson, D.T. (593) 128
Reisert, I., see Beyer, C. (593) 253
Ren, C., see Shen, E. (593) 136
Rhoads, D.E., see Foley, T.D. (593) 39
Richey, P., see Tabaton, M. (593) 299
Rittenhouse, P.A., Li, Q., Levy, A.D. and Van de Kar, L.D.
Neurons in the hypothalamic paraventricular nucleus mediate the serotonergic stimulation of renin secretion (593) 105
Roberts, B.L.
Differences in the dopaminergic innervation of the electroreceptive and mechanoreceptive medullary lateral line nuclei of the ray, *Raja radiata* (593) 339
Robertson, H.A., Peterson, M.R. and Worth, G.G.
Synergistic and persistent interaction between the D2 agonist, bromocriptine, and the D1 selective agonist, CY 208–243 (593) 332
Robinson, P.P., see Doubleday, B. (593) 179
Rossini, P.M., Desiato, M.T. and Caramia, M.D.
Age-related changes of motor evoked potentials in healthy humans: Non-invasive evaluation of central and peripheral motor tracts excitability and conductivity (593) 14

Saito, H., see Abe, K. (593) 335
Saito, H., see Chida, N. (593) 57
Saitoh, T., see Masliah, E. (593) 323
Sakurada, S., see Sakurada, T. (593) 319
Sakurada, T., Manome, Y., Tan-No, K., Sakurada, S., Kisara, K., Ohba, M. and Terenius, L.
A selective and extremely potent antagonist of the neurokinin-1 receptor (593) 319
Serratosa, J., see Tusell, J.M. (593) 209
Shen, E., Dun, S.L., Ren, C., Bennett-Clarke, C. and Dun, N.J.
Hypotension preferentially induces *C-fos* immunoreactivity in supraoptic vasopressin neurons (593) 136
Shigematsu, K. and McGeer, P.L.
Accumulation of amyloid precursor protein in damaged neuronal processes and microglia following intracerebral administration of aluminum salts (593) 117
Sigworth, S.K., see Hyde, T.M. (593) 291

Simon, R.P., see Meldrum, B.S. (593) 1
Stauch Slusher, B., see Meyerhoff, J.L. (593) 140
Steinberg, R., see Lafon-Cazal, M. (593) 63
Stephenson, D.T., Rash, K. and Clemens, J.A.
Amyloid precursor protein accumulates in regions of neurodegeneration following focal cerebral ischemia in the rat (593) 128
Stratta, F., see Mercuri, N.B. (593) 51
Sugiura, H. and Yamauchi, T.
Developmental changes in the levels of Ca^{2+}/calmodulin-dependent protein kinase II α and β proteins in soluble and particulate fractions of the rat brain (593) 97
Sutula, T.P., see Golarai, G. (593) 257
Swan, J.H., see Meldrum, B.S. (593) 1

Tabaton, M., Cammarata, S., Mandybur, T., Richey, P., Kawai, M., Perry, G. and Gambetti, P.
Senile plaques in cerebral amyloid angiopathy show accumulation of amyloid precursor protein without cytoskeletal abnormalities (593) 299
Tan-No, K., see Sakurada, T. (593) 319
Terenius, L., see Sakurada, T. (593) 319
Thomas, I., see Chance, W.T. (593) 20
Tooyama, I., Walker, D., Yamada, T., Hanai, K., Kimura, H., McGeer, E.G. and McGeer, P.L.
High molecular weight basic fibroblast growth factor-like protein is localized to a subpopulation of mesencephalic dopaminergic neurons in the rat brain (593) 274
Tortorella, A., see Pasini, A. (593) 287
Trullas, R., see Tusell, J.M. (593) 209
Tusell, J.M., Vendrell, M., Serratosa, J. and Trullas, R.
Lindane-induced convulsions in NMRI and OF1 mice: antagonism with (+)MK-801 and voltage-dependent calcium channel blockers (593) 209

Van de Kar, L.D., see Rittenhouse, P.A. (593) 105
Vanderwolf, C.H.
Hippocampal activity, olfaction, and sniffing: an olfactory input to the dentate gyrus (593) 197
Vendrell, M., see Tusell, J.M. (593) 209

Walker, D., see Tooyama, I. (593) 274
Walker, J.M., see Hohmann, A.G. (593) 265
Weiss, F., Markou, A., Lorang, M.T. and Koob, G.F.
Basal extracellular dopamine levels in the nucleus accumbens are decreased during cocaine withdrawal after unlimited-access self-administration (593) 314
Widmer, H., Dreifuss, J.J. and Raggenbass, M.
N-Methyl-D-aspartate and vasopressin activate distinct voltage-dependent inward currents in facial motoneurones (593) 215
Wiechmann, A.F. and Wirsig-Wiechmann, C.R.
Asymmetric distribution of melatonin receptors in the brain of the lizard *Anolis carolinensis* (593) 281
Wirsig-Wiechmann, C.R., see Wiechmann, A.F. (593) 281
Worth, G.G., see Robertson, H.A. (593) 332
Wu, L.-C., see Hyde, T.M. (593) 291

Yamada, T., see Tooyama, I. (593) 274
Yamauchi, T., see Sugiura, H. (593) 97
Yim, G.K.W., see Patel, M.N. (593) 114
Yourick, D.L., see Meyerhoff, J.L. (593) 140

Zhang, D.X. and Levy, W.B.
Ketamine blocks the induction of LTP at the lateral entorhinal cortex–dentate gyrus synapses (593) 124
Zvartau, E., see Kuzmin, A. (593) 221